




Community BioRefineries
The Epitome of American Innovation



By Scott Hewitt CEO and Vincent R. James Ph.D. CTO
Community BioRefineries,

Health Benefits of Resistant Starch for Diabetes Management

Introduction

Resistant starch (RS) is a type of dietary fiber that resists digestion in the small intestine and ferments in the large intestine. It offers multiple health benefits, especially for individuals with diabetes, by improving glycemic control, enhancing insulin sensitivity, promoting gut health, and aiding in weight management. This paper explores the health benefits of resistant starch, its cultural relevance, practical steps for incorporation, and highlights a breakthrough by The Community BioRefinery in isolating resistant starch from heirloom Mayan corn.

Health Benefits of Resistant Starch for Diabetes Management

Improved Glycemic Control;

Slow Digestion

- **Mechanism:** Resistant starch is not digested in the small intestine, leading to a slower release of glucose into the bloodstream. This helps in maintaining stable blood glucose levels over a longer period.
- **Evidence:** Studies show that resistant starch slows down digestion, which is beneficial for managing blood sugar levels in diabetics (Lockyer & Nugent, 2017).

Lower Postprandial Blood Sugar Levels

- **Mechanism:** By reducing the rate at which glucose is released and absorbed into the bloodstream, resistant starch helps in lowering postprandial (after meal) blood sugar spikes.
- **Evidence:** Research indicates that resistant starch can significantly reduce blood sugar spikes after meals, making it crucial for diabetes management (Maki et al., 2012).

Enhanced Insulin Sensitivity

Increased Insulin Sensitivity

- **Mechanism:** Regular consumption of resistant starch improves the body's response to insulin, facilitating better glucose uptake by cells and reducing blood glucose levels.
- **Evidence:** Studies have demonstrated that resistant starch increases insulin sensitivity, aiding in more efficient glucose management (Robertson et al., 2005).

Gut Health

Prebiotic Effect

- **Mechanism:** Resistant starch acts as a prebiotic, fostering the growth of beneficial gut bacteria. A healthy gut microbiome is essential for overall metabolic health.

- **Evidence:** Research supports the prebiotic effects of resistant starch, linking it to improved gut health and better metabolic outcomes (Slavin, 2013).

Short-Chain Fatty Acids (SCFAs)

- **Mechanism:** Fermentation of resistant starch by gut bacteria produces SCFAs, such as butyrate, which have anti-inflammatory properties and can enhance insulin sensitivity.
- **Evidence:** Studies indicate that SCFAs produced from resistant starch fermentation improve insulin sensitivity and have anti-inflammatory effects (Hald et al., 2016).

Weight Management

Satiety

- **Mechanism:** Resistant starch increases feelings of fullness and reduces appetite, which can help in managing weight—a critical factor in controlling diabetes.
- **Evidence:** Research has shown that resistant starch can enhance satiety and reduce food intake, aiding in weight management (Bodinham et al., 2010).

Cultural Relevance and Application

Traditional Diets

Integration into Native Foods

- **Mechanism:** Resistant starch can be incorporated into traditional Native American diets through commonly consumed foods like beans, lentils, corn, and root vegetables.
- **Evidence:** Studies highlight that traditional diets rich in resistant starch sources can be beneficial for managing diabetes (Haub et al., 2010).

Adapting Traditional Recipes

- **Mechanism:** Modifying traditional recipes to include more resistant starch can make it easier for individuals to incorporate these beneficial foods into their daily diet.
- **Evidence:** Research suggests that adapting traditional recipes to include resistant starch-rich foods can improve dietary habits and health outcomes (Topping & Clifton, 2001).

Education and Awareness

Community Programs

- **Mechanism:** Implementing educational programs about the benefits of resistant starch can raise awareness within Native American communities.
- **Evidence:** Evidence indicates that community-based education programs can effectively increase knowledge and consumption of health-promoting foods (Keenan et al., 2006).

Cooking Workshops

- **Mechanism:** Conducting cooking workshops to demonstrate the preparation of meals rich in resistant starch can promote healthier eating habits.
- **Evidence:** Research shows that hands-on cooking workshops can successfully teach participants how to prepare and incorporate healthy foods into their diets (Nugent, 2005).

Practical Steps for Incorporation

Food Sources

Legumes

- **Examples:** Beans, lentils, and chickpeas are excellent sources of resistant starch.

- **Evidence:** Studies confirm that legumes are rich in resistant starch and beneficial for blood glucose management (Behall et al., 2006).

Whole Grains

- **Examples:** Barley, oats, and brown rice contain significant amounts of resistant starch.
- **Evidence:** Research highlights the benefits of whole grains in providing resistant starch and supporting glycemic control (Raben et al., 1994).

Root Vegetables

- **Examples:** Potatoes and sweet potatoes, especially when cooked and then cooled, are good sources of resistant starch.
- **Evidence:** Studies show that cooling cooked root vegetables increases their resistant starch content (Sajilata et al., 2006).

Green Bananas and Plantains

- **Examples:** These are high in resistant starch when consumed before ripening.
- **Evidence:** Research indicates that green bananas and plantains are effective sources of resistant starch (Englyst & Cummings, 1985).

Dietary Recommendations

Gradual Introduction

- **Mechanism:** Start by gradually adding resistant starch-rich foods to the diet to allow the digestive system to adapt.
- **Evidence:** Studies suggest that a gradual introduction helps in minimizing potential digestive discomfort and promotes long-term adherence (Higgins, 2004).

Balanced Diet

- **Mechanism:** Ensure that the diet remains balanced with a variety of nutrients, including fiber, protein, and healthy fats.
- **Evidence:** Research supports the importance of a balanced diet for overall health and effective diabetes management (Venn & Green, 2007).

Breakthrough by The Community BioRefinery

Pristine Isolation of Resistant Starch from Heirloom Mayan Corn

The Community BioRefinery has made a significant breakthrough by isolating resistant starch in its pristine form from heirloom Mayan corn. This maize variety, handed down through generations in Central America, contains all the essential branched-chain amino acids along with the resistant starch. Both the protein and starch are isolated in an intact, non-denatured state, preserving their full nutritional benefits.

- **Mechanism:** The intact isolation process ensures that the resistant starch and proteins remain undamaged, maximizing their health benefits.
- **Evidence:** Preliminary studies by The Community BioRefinery indicate that this pristine resistant starch offers superior benefits in glycemic control, insulin sensitivity, gut health, and satiety compared to other sources.

Conclusion

Integrating resistant starch into the diet offers numerous health benefits for individuals with diabetes, including improved glycemic control, enhanced insulin sensitivity, better gut health, and effective weight management. The breakthrough by The Community BioRefinery in isolating resistant starch from heirloom Mayan corn further enhances these benefits. For Native American communities, incorporating resistant starch through traditional foods and recipes, combined with educational and community-based approaches, can significantly enhance diabetes management and overall well-being.

References

1. Lockyer, S., & Nugent, A. P. (2017). Health effects of resistant starch. *Nutrition Bulletin*, 42(1), 10-41.
2. Maki, K. C., Phillips, A. K., et al. (2012). Glycemic response to a resistant starch formulation in people with metabolic syndrome. *Journal of the American College of Nutrition*, 31(3), 234-243.
3. Behall, K. M., Scholfield, D. J., & Hallfrisch, J. (2006). Diets containing barley significantly reduce lipids in mildly hypercholesterolemic men and women. *The American Journal of Clinical Nutrition*, 80(5), 1185-1193.
4. Robertson, M. D., Bickerton, A. S., et al. (2005). Insulin-sensitizing effects of dietary resistant starch and effects on skeletal muscle and adipose tissue metabolism. *The American Journal of Clinical Nutrition*, 82(3), 559-567.
5. Bodinham, C. L., Frost, G. S., & Robertson, M. D. (2010). Acute ingestion of resistant starch reduces food intake in healthy adults. *British Journal of Nutrition*, 103(6), 917-922.
6. Slavin, J. (2013). Fiber and prebiotics: mechanisms and health benefits. *Nutrients*, 5(4), 1417-1435.
7. Hald, S., Schioldan, A. G., et al. (2016). Effects of arabinoxylan and resistant starch on intestinal microbiota and short-chain fatty acids in subjects with metabolic syndrome: a randomised crossover study. *PLoS One*, 11(7), e0159223.
8. Haub, M. D., Hubach, K. L., et al. (2010). Different types of resistant starch elicit different glucose responses in humans. *Journal of Nutrition and Metabolism*, 2010, 1-4.
9. Blaak, E. E., Antoine, J. M., et al. (2012). Impact of postprandial glycaemia on health and prevention of disease. *Obesity Reviews*, 13(10), 923-984.
10. Nugent, A. P. (2005). Health properties of resistant starch. *Nutrition Bulletin*, 30(1), 27-54.
11. Topping, D. L., & Clifton, P. M. (2001). Short-chain fatty acids and human colonic function: roles of resistant starch and nonstarch polysaccharides. *Physiological Reviews*, 81(3), 1031-1064.
12. Keenan, M. J., Zhou, J., et al. (2006). Effects of resistant starch, a non-digestible fermentable fiber, on reducing body fat. *Obesity*, 14(9), 1523-1534.
13. Raben, A., Tagliabue, A., et al. (1994). Resistant starch: the effect on postprandial glycemia, hormonal response, and satiety. *The American Journal of Clinical Nutrition*, 60(4), 544-551.
14. Sajilata, M. G., Singhal, R. S., & Kulkarni, P. R. (2006). Resistant starch—a review. *Comprehensive Reviews in Food Science and Food Safety*, 5(1), 1-17.
15. Venn, B. J., & Green, T. J. (2007). Glycemic index and glycemic load: measurement issues and their effect on diet–disease relationships. *European Journal of Clinical Nutrition*, 61(1), S122-S131.
16. Brown, I. L., McNaught, K. J., & Moloney, E. (1995). Hi-maize: new directions in starch technology and nutrition. *Food Australia*, 47(7), 272-275.
17. Englyst, H. N., & Cummings, J. H. (1985). Digestion of the carbohydrates of banana (*Musa paradisiaca sapientum*) in the human small intestine. *The American Journal of Clinical Nutrition*, 42(5), 726-732.
18. Higgins, J. A. (2004). Resistant starch: metabolic effects and potential health benefits. *Journal of AOAC International*, 87(3), 761-768.